

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|--|
| (51) International Patent Classification ⁶ : C07D 209/08, A61K 31/40, C07D 209/30, 409/12, 209/12 | | A1 | (11) International Publication Number: WO 98/08817 |
| | | | (43) International Publication Date: 5 March 1998 (05.03.98) |
| (21) International Application Number: PCT/US97/15026 (22) International Filing Date: 26 August 1997 (26.08.97) (30) Priority Data: 08/703,562 27 August 1996 (27.08.96) US (71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US). (72) Inventors: MEWSHAW, Richard, Eric; 21 Boxwood Drive, Princeton, NJ 08540 (US). WEBB, Michael, Byron; Apartment 2401, 9071 Mill Creek Road, Levittown, PA 19054 (US). (74) Agents: ALICE, Ronald, W.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US) et al. | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> | |
| (54) Title: 4-AMINOETHOXY INDOLES AS DOPAMIN D2 AGONISTS AND AS 5HT _{1A} LIGANDS | | | |
| (57) Abstract Compounds of the formula (I) in which R ₁ is hydrogen, alkyl, cycloalkylalkyl, arylalkyl, (haloaryl)alkyl, (alkoxy-aryl)alkyl, thienylmethyl, furanylmethyl, pyridinylmethyl, alkylphenyl, 4-fluoro-butylphenyl or 6-fluoro-1,2-benzisoxazol-yl-propyl; X is hydrogen, halogen, cyano, alkyl, acetyl, trifluoroacetyl, trifluoromethyl or formyl; Y is hydrogen, halogen, alkoxy or alkyl; or a pharmaceutically acceptable salt thereof are inhibitors of dopamine synthesis and release, useful in the treatment of schizophrenia, Parkinson's Disease, Tourette's Syndrome, alcohol addiction, cocaine addiction, and addiction to analogous drugs and they also have affinity for the 5-HT _{1A} receptors which characterizes them as useful in the treatment of diseases attending disturbances in the serotonergic systems, such as anxiety, stress, depression, sexual dysfunctions and sleep disturbances. | | | |
| | | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

- 1 -

4-AMINOETHOXY INDOLES AS DOPAMINE D₂ AGONISTS AND AS 5HT_{1A} LIGANDS**BACKGROUND OF INVENTION**

5 Efforts to induce antipsychotic activity with dopamine autoreceptor agonists have been successful [Dorsini et al., Adv. Biochem. Psychopharmacol 16, 645-648, (1977); Tamminga et al., Science 200, 567-568; and Tamminga et al., Psychiatry 398-402, (1986)]. A method for determining intrinsic activity at the dopamine D₂ receptor was recently reported [Lahti et al., Mol. Pharm. 42, 432-438, (1993)].

10 Intrinsic activity is predicted using the ratio of the "low-affinity agonist" (LowAg) state of the receptor and the "high-affinity agonist" (HighAg) state of the receptor, i.e. LowAg/HighAg. These ratios correlate with the agonist, partial agonist and antagonist activities of a given compound, which activities characterize a compound's ability to elicit an antipsychotic effect.


15 U.S. Pat. Nos. 3,906,000 and 3,904,645 describe a series of indoles which are useful as oral hypoglycemic agents. Troxler et al. 66-25558F: WPIDS describes a series of indoles including 4-(2-hydroxy-3-isopropyl- or secondary butyl-amino-propoxy)-indoles which are useful as β -adrenergic blocking agents for the treatment of

20 heart diseases.

DESCRIPTION OF THE INVENTION

25 In accordance with this invention, there is provided a group of aminoethoxy indole derivatives which are useful antipsychotic agents. In addition, this invention provides processes for preparation of the compounds and methods for their use in treating diseases of the central nervous system. The aminoethoxy indoles of this invention are illustrated by the following Formula I:

- 2 -



I

in which:


R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cycloalkylalkyl of 6 to 12 carbon atoms, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms, (alkoxyaryl)alkyl of 8 to 12 carbon atoms, thienylmethyl, furanylmethyl, pyridinylmethyl, alkylphenyl of 7 to 12 carbon atoms, 4-fluorobutyrophenone or 6-fluoro-1,2-benzisoxazol-yl-propyl;

X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.

More specifically, the compounds of this invention are 4-aminoethoxy-indoles illustrated by Formula I.



I

in which:

- 3 -

R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cyclohexylmethyl, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms or (alkoxyaryl)alkyl of 8 to 12 carbon atoms;

5 X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.


10 More preferred compounds are those of Formula I in which R₁ is alkyl of 1 to 6 carbon atoms, benzyl, halobenzyl, alkoxybenzyl of 8 to 12 carbon atoms or alkylbenzyl of 8 to 12 carbon atoms; X is hydrogen, halogen or trifluoroacetyl and Y is hydrogen or halogen; or a pharmaceutically acceptable salt thereof.

15 The pharmaceutically acceptable acids from which addition salts are conventionally produced, having the same utility as the free base, include both inorganic or organic acids. For example: fumaric, maleic, benzoic, ascorbic, pantoic, succinic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic,
20 citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene-sulfonic, hydrochloric hydrobromic, sulfuric, cyclohexylsulfamic, phosphoric, nitric acid, and the like, are suitable for this purpose.

25 The compounds of Formula I are generally prepared by the overall reaction sequence indicated in Schemes I, II and III as follows:


- 4 -

Scheme I





- 5 -

Scheme II



- 6 -

Route III**Scheme III**

5

The compounds of this invention are dopamine agonists with various degrees of intrinsic activity. Some are selective autoreceptor agonists and others bind to the postsynaptic D₂ receptors. The autoreceptor agonists act as partial agonists (i.e. activate only autoreceptors versus postsynaptic D₂ dopamine receptors). As such, they provide functional modulation of the dopamine systems of the brain without the

10

- 7 -

excessive blockade of the postsynaptic dopamine receptors which have been observed to be responsible for the serious side effects frequently exhibited by agents found otherwise clinically effective for the treatment of schizophrenia. Activation of the dopamine autoreceptors results in reduced neuronal firing as well as inhibition of dopamine synthesis and release and therefore provide a means of controlling hyperactivity of the dopaminergic systems with essentially no extrapyramidal side effects (EPS).

The compounds of this invention were also found to have affinity for the 5-HT_{1A} receptors and therefore have the ability to modulate serotonergic activity. As such, they are useful in the treatment of diseases characterized by disturbances in the dopaminergic and serotonergic systems, such as schizophrenia, Parkinson's disease, Tourette's Syndrome, alcohol addiction, cocaine addiction, anxiety, stress, depression, sexual dysfunctions and sleep disturbances.

The following examples illustrate, without limitation, methods for production of the compounds of this invention.

Intermediate 1

N-Benzyl-N-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester

A solution of N-benzylaminoethanol (4.8 g, 31.9 mmol) and di-tertbutyl-dicarbonate (7.5 g, 34.4 mmol) in anhydrous tetrahydrofuran (30 mL) was stirred at ambient temperature for 18 hours. The solvent was removed and the product purified by flash chromatography (ethyl acetate-hexane, 1:1) to afford 8.0 g (99%) of a thick oil.

Elemental analysis for C₁₄H₂₁NO₃

Calc'd: C, 66.91; H, 8.42; N, 5.57

Found: C, 66.64; H, 8.59; N, 5.60

This general procedure utilizing N-methylaminoethanol afforded:

(1b) N-Methyl-N-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester as a clear oil (83.7%); MS m/z 175 (M⁺).

- 8 -

Elemental analysis for $C_8H_{17}NO_3$

Calc'd C, 54.84; H, 9.78; N, 7.99

Found C, 54.35; H, 10.00; N, 7.84

5

Intermediate 2

Method A

N-Benzyl-N-[2-(1H-indol-4-yloxy)-ethyl]- carbamic acid tert-butyl ester

10

To a solution of benzyl-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester (12.68 g, 50.5 mmol), 4-hydroxyindole (4.48 g, 33.6 mmol) and triphenylphosphine (14.1 g, 53.8 mmol) in anhydrous tetrahydrofuran (130 mL) was slowly added a solution of diethylazidocarboxylate (9.38 g, 53.8 mmol) in tetrahydrofuran (15 mL) at room temperature. The reaction mixture was stirred for 16 hours and then the solvent was removed and the crude product dissolved in diethyl ether and diluted with hexanes. After standing for 30 minutes, the solid was filtered and the filtrate concentrated. The product was purified by flash chromatography to afford 8.6 g of a yellow oil (69.7 %).

15

Elemental analysis for $C_{22}H_{26}N_2O_3$

Calc'd: C, 72.11; H, 7.15; N, 7.64

Found: C, 71.39; H, 7.28; N, 7.21

This general procedure utilizing N-methyl-N-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester, N-(2-hydroxyethyl)-phthalimide afforded, chloroethanol or 4-chloro-3-nitrophenol afforded, respectively:

25

(2b) N-Methyl-N-[2-(1H-indol-4-yloxy)-ethyl]-carbamic acid tert-butyl ester as a yellow oil; (77.2 %); MS EI m/z 290 (M+).

30

(2c) N-[2-(1H-Indol-4-yloxy)-ethyl]-phthalimide as a white solid: (13.2 %); mp 155-157°C; IR (KBr) 3400, 1725 cm^{-1} ; MS EI m/e 306 (M+).

- 9 -

Elemental analysis for $C_{18}H_{14}N_2O_3$

Calc'd: C, 70.58; H, 4.61; N, 9.15.

Found: C, 70.33; H, 4.43; N, 9.11

5 (2d) 2-(1H-Indol-4-yloxy)-chloroethane: (57 %), mp 62-63 °C.

(2e) 1-(2-Chloroethoxy)-4-chloro-3-nitrobenzene: (93%); mp 46-48 °C; MS EI *m/e* 235, 237, 239 (M^+); 1H NMR (400 MHz, DMSO- d_6) δ 3.95 (t, 2H, $J=5.2$ Hz), 4.36 (t, 2H, $J=5.2$ Hz), 7.32 (dd, 1H, $J=3.2$, $J=8.9$ Hz), 7.66, (d, 1H, $J=9$ Hz), 7.69, (d, 1H, $J=3.2$ Hz).

10

Elemental analysis for $C_8H_7Cl_2NO_3$

Calc'd: C, 40.71; H, 2.99; N, 5.93.

Found: C, 40.43, H, 2.71; N, 5.62.

15

(2f) 1-(2-Chloroethoxy)-4-chloro-3-nitrobenzeneMethod B

To a 2L 3-neck round-bottom flask was added 4-chloro-3-nitro-phenol (50 g, 0.29 mol), potassium carbonate (100g, 0.72 mol), dichloroethane (315 g, 3.2 mol), potassium iodide (5 g) and 2-butanone (1 L). The mixture was mechanically stirred and heated to reflux for 44 hours then allowed to cool to room temperature and the solids were filtered. The solvent was evaporated under vacuum and the oil dissolved in diethyl ether (300 mL) and washed with 10 % sodium hydroxide. The organic layer was dried over anhydrous magnesium sulfate, filtered, and the solvent removed under vacuum. The product was dissolved in 1:1 methylene chloride-hexanes and filtered through silica. Upon concentration 54.5 g (78. % %) of product was afforded as a white solid: mp 44.5-46 °C.

25

30 Elemental analysis for $C_8H_7Cl_2NO_3$

Calc'd: C, 40.71; H, 2.99; N, 5.93.

Found: C, 40.89, H, 2.70; N, 5.83.

- 10 -

Intermediate 3**7-Chloro-4-(2-chloroethoxy)-1H-indole**

To a solution of 1-(2-chloroethoxy)-4-chloro-3-nitrobenzene (10.00 g, 0.04236 mol) in THF (230 mL) stirred in a cold bath at -50 to -40 °C was added a THF solution of vinylmagnesium bromide (132 mL, 1.0 M, 0.132 mol) over 2 minutes. After stirring in the cold bath for 2-2.5 hours, saturated NH₄Cl (150 mL) was added to the cold solution and it was removed from the cold bath. Enough 1 M HCl was added to dissolve the precipitated solids. This two phase system was stirred for 0.5 hour at most. The layers were separated and the aqueous phase was extracted once with Et₂O. Combination of the Et₂O and THF followed by drying over MgSO₄ and evaporation gave 15.43 g of a dark oil. This was purified by chromatography on silica gel using a three component elutant which consisted of 80% CH₂Cl₂ and 20% of a gradient of EtOAc/hexane. This gave the product as a yellow solid: 3.32 g (34%); mp 68-72 °C; MS EI *m/e* 229, 231, 233 (M⁺); ¹H NMR (400 MHz, DMSO-d₆) δ 3.99 (t, 2H, J=5.1 Hz), 4.34 (t, 2H, J=5.0 Hz), 6.51 (t, 1H, J=2.7 Hz), 6.53 (d, 1H, J=7.8 Hz), 7.04 (d, 1H, J=8.0 Hz), 7.29 (t, 1H, J=2.7 Hz), 11.43 (s, 1H).

Elemental analysis for C₁₀H₉Cl₂NO

Calc'd: C, 52.20; H, 3.94; N, 6.09.

Found: C, 52.09; H, 3.92; N, 5.96.

Intermediate 4**3,7-Dichloro-4-(2-chloroethoxy)-1H-indole**

To a solution of 7-chloro-4-(2-chloroethoxy)-1H-indole (4.61 g, 20.0 mmol) in acetonitrile (100 mL) was added N-chlorosuccimide (2.94 g, 2.20 mmol) at room temperature. The reaction was allowed to stir for 1.5 hour then poured into water (100 mL) and extracted with methylene chloride (200 mL). The organic layer was dried over anhydrous magnesium sulfate, filtered, and the solvent removed under vacuum to afford a dark solid. This material was chromatographed (methylene chloride-hexanes: 1:2) to afford 4.15 g (78.4 %) as a white solid: mp 106-107.5 °C; IR (KBr) 3400 cm⁻¹; MS EI *m/e* 263, 265, 267, 269 (M⁺); ¹H NMR (CDCl₃) δ 3.91 (2H, t, J=6.2 Hz), 4.33 (2H, t, J=6.2 Hz), 6.47 (1H, d, J= 8.4 Hz), 7.08-7.13 (2H, m), 8.26 (1H, bs, NH).

- 11 -

Elemental analysis for $C_{22}H_{25}N_2O_3Cl$

Calc'd: C, 65.91; H, 6.28; N, 6.99

Found: C, 65.61; H, 6.21; N, 6.89

5

EXAMPLE 1**[2-(1H-Indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-amine**

A solution of the 2-(1H-indol-4-yloxy)-chloroethane (1.80 g, 9.20 mmol) and
10 4-phenyl-1-aminobutane (4.12g, 27.6 mmol) in anhydrous dimethylsulfoxide (25 mL)
was heated to 80 °C for 6 hours. The reaction mixture was poured into water (150 mL)
and extracted with methylene chloride (3x 100 mL). The organic layers were combined
and dried over anhydrous magnesium sulfate, filtered, and the solvent concentrated.
Purification by flash chromatography (5%-10% methanol-CH₂Cl₂) afforded 1.89 g
15 (65.9%) of a tan oil: MS *m/e* 308 (M⁺). The oxalate salt was prepared in
tetrahydrofuran: mp 202-204 °C.

Elemental analysis for $C_{20}H_{24}N_2O \cdot C_2H_2O_4 \cdot 0.5H_2O$

Calc'd: C, 64.85; H, 6.68; N, 6.87.

20 Found: C, 64.66; H, 6.61; N, 6.70.

This general procedure utilizing 7-chloro-4-(2-chloroethoxy)-1H-indole or 3,7-
dichloro-4-(2-chloroethoxy)-1H-indole and reacting with either benzylamine; 4-
fluorobenzyl amine, 4-chlorobenzyl amine or thiophene-2-methylamine afforded:

25

(1b) Benzyl-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-amine (68%). The fumarate salt
was prepared in isopropanol as colorless crystals; mp 168-170 °C; MS EI *m/e* 300, 302
(M⁺).

30 Elemental analysis for $C_{17}H_{17}ClN_2O \cdot 0.5C_4H_4O_4 \cdot 0.25C_3H_8O$

Calc'd: C, 63.45; H, 5.66; N, 7.49

Found: C, 63.12; H, 5.61; N, 7.31.

- 12 -

(1c) Benzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine (67.8 %): The fumarate salt was prepared and characterized: mp 201-202 °C; MS EI *m/e* 334, 336, 338 (M^+).

5 Elemental analysis for $C_{17}H_{16}Cl_2N_2O \cdot 0.5C_4H_4O_4$
 Calc'd: C, 58.03; H, 4.61; N, 7.12.
 Found: C, 57.88; H, 4.45; N, 6.96.

(1d) 4-Fluorobenzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine (64.5 %): mp
10 102.5-103.5 °C.

 Elemental analysis for $C_{17}H_{15}FCl_2N_2O$
 Calc'd: C, 57.81; H, 4.28; N, 7.93.
 Found: C, 57.68; H, 4.16; N, 7.86.

15 (1e) 4-Chlorobenzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine (59.9 %): mp
 115-116 °C; MS EI 368 *m/e* (M^+).

 Elemental analysis for $C_{17}H_{15}Cl_3N_2O \cdot 0.25H_2O$
20 Calc'd: C, 54.57; H, 4.17 ; N, 7.49.
 Found: C, 54.43; H, 3.82; N, 7.32.

(1f) Thien-2-ylmethyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine (76.3%):
mp 99-101 °C, MS EI 340, 342, 344 *m/e* (M^+).

25 Elemental analysis for $C_{15}H_{15}Cl_2N_2OS$
 Calc'd: C, 52.70; H, 4.13 ; N, 8.21.
 Found: C, 52.70; H, 3.95; N, 8.19.

30 **Intermediate 5**
 N-Benzyl-N-[2-(1H-indol-3-(2,2,2-trifluoroethanoyl)-4-
 yloxy)-ethyl]-carbamic acid tert-butyl ester

 To a stirring anhydrous solution of benzyl-[2-(1H-indol-4-yloxy)-ethyl]-
35 carbamic acid tert-butyl ester (1.85 g, 5.05 mmol) and TEA (0.8 mL, 0.6 g, 6 mmol) in

- 13 -

CH₂Cl₂ was added trifluoroacetic acid anhydride (1.1 mL, 1.6 g, 7.8 mmol) over 5 minutes at room temperature. The reaction mixture was stirred at room temperature over-night. It was washed twice with H₂O and then dried over MgSO₄. Evaporation of solvent gave 3.38g of residue. This was purified by chromatography on silica gel with a hexane/EtOAc gradient to give the title compound as an amorphous light yellow solid: 1.15g (49%); MS EI *m/e* 462 (M⁺); IR(KBr) 1719 cm⁻¹, 1744 cm⁻¹; ¹H NMR (400 MHz, DMSO-d₆) δ 1.33 and 1.38 (2s, 9H, rotamers), 3.50-3.63 (2m, 2H, rotamers), 4.15 (t, 2H, J=5.5 Hz), 4.54 (s, 2H), 6.77 (d, 1H, J=7.9 Hz), 7.15 (d, 1H, J=8.1 Hz), 7.20-7.27 (m, 4H), 7.29-7.35 (m, 2H), 8.32 (s, 1H), 12.58 (s, 1H).

Intermediate 6

N-[2-(1H-indol-4-yloxy)-ethyl]-N-(4-phenyl-butyl)-trifluoroacetamide

To a solution of [2-(1H-indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-amine (2.38 g, 7.72 mmol) and triethylamine (1.56 g, 15.4 mmol) in anhydrous methylene chloride (30 mL) at room temperature was slowly added trifluoroacetic anhydride (2.42 g, 11.6 mmol) over 10 minutes. The reaction was stirred for 1 hour and then poured into a 1:1 solution of saturated sodium carbonate-water (50 mL) and extracted with methylene chloride (2x100 mL). The organic layer dried over anhydrous magnesium sulfate, filtered, and the solvent evaporated. Purification by flash chromatography (20% ethyl acetate-hexanes) afforded 1.61 g (51.6%) of an off-white solid: mp 70-72 °C, MS *m/e* 404 (M⁺); IR (KBr) 3360, 2950, 1725 cm⁻¹.

Intermediates 7 & 8

N-Benzyl-N-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide

and

N-Benzyl-N-[2-(7-chloro-3-trifluoroacetyl-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide

To a solution of benzyl-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-amine (4.55 g, 15.1 mmol) in CH₂Cl₂ (200 mL) at room temperature was added triethylamine (TEA) (2.15 mL, 1.56 g, 15.4 mmol) and then trifluoroacetic acid anhydride (4.5 mL, 6.7 g, 32 mmol) over 20 minutes. The solution was stirred at room temperature over-night. It

- 14 -

was washed twice with H₂O. Drying over MgSO₄ and evaporation of the solvent gave 7.33 g of residue which consisted primarily of the two products. These were separated and purified by chromatography on silica gel with a gradient of CH₂Cl₂/hexane/EtOAc (10/80/10, 4/82/14, 0/86/14, 0/80/20) which first eluted N-benzyl-N-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide as light yellow crystals: 2.79 g (47%); mp 114-116 °C; MS EI *m/e* 396 (M⁺); IR (KBr) 1682 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 3.79 and 3.86 (2t, 2H, J=5.6 Hz, J=5.0 Hz, rotamers), 4.28 and 4.31 (2t, 2H, J=5.6 Hz, J=5.0 Hz, rotamers), 4.89 and 4.93 (2s, 2H, rotamers), 6.38 and 6.40 (2d, 1H, J=8.3 Hz, J=8.5 Hz, rotamers), 6.64-6.68 (m, 1H), 7.05 and 7.08 (2d, 1H, J=8.1 Hz, J=8.3 Hz, rotamers); 7.19-7.44 (m, 6H), 8.42 (s, 1H).

Elemental analysis for C₁₉H₁₆ClF₃N₂O₂

Calc'd: C, 57.51; H, 4.06; N, 7.06.

Found: C, 57.11; H, 3.88; N, 7.01.

15

N-Benzyl-N-[2-(7-chloro-3-trifluoroacetyl-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide was then eluted off the column to afford 3.18 g (43%) of crystalline solid; mp 152-154 °C; MS FAB *m/e* 493 (MH⁺); IR (KBr) 1685 cm⁻¹, 1699 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 3.84 and 4.01 (2t, 2H, J=5.0 Hz, J=5.3 Hz, rotamers), 4.26 and 4.31 (2t, 2H, J=5.5 Hz, J=5.0 Hz, rotamers), 4.92 and 5.00 (2s, 2H, rotamers), 6.63 and 6.66 (2d, 1H, J=8.8 Hz, J=8.8 Hz, rotamers), 7.27-7.42 (m, 6H), 8.04-8.08 (m, 1H), 9.13 (s, 1H).

20

Elemental analysis for C₂₁H₁₅ClF₆N₂O₃)

25

Calcd: C, 51.18; H, 3.07; N, 5.68.

Found: C, 51.31; H, 2.89; N, 5.58.

Intermediate 9

N-Benzyl-N-[2-(3-chloro-1H-indol-4-yloxy)-ethyl]- carbamic acid tert-butyl ester

30

To a solution of N-benzyl-N-[2-(1H-indol-4-yloxy)-ethyl]-carbamic acid tert-butyl ester (6.3 g, 17.2 mmol) in tetrahydrofuran (100 mL) was added N-chlorosuccinimide (2.3 g, 17.2 mmol) in two portions over 1 hour. The reaction was allowed to stir for 18 hours and the solvent removed under vacuum. The mixture was

35

- 15 -

dissolved in diethyl ether and the insoluble solids filtered. The solvent was again removed and the product purified by chromatography (30% ethyl acetate-hexanes) to afford 5.65 g of white solid (81.9 %): mp 114-116 °C.

5 Elemental analysis for $C_{22}H_{25}N_2O_3Cl$

Calc'd: C, 65.91; H, 6.28; N, 6.99

Found: C, 65.61; H, 6.21; N, 6.89

10 This general procedure utilizing N-methyl-N-[2-(1H-indol-4-yloxy)-ethyl]-carbamic acid tert-butyl ester, N-[2-(1H-Indol-4-yloxy)-ethyl]-phthalimide, N-benzyl-N-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide and [2-(1H-indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-trifluoroacetamide afforded, respectively:

15 (9b) N-Methyl-N-[2-(3-chloro-1H-indol-4-yloxy)-ethyl]-carbamic acid tert-butyl ester as a white solid: (74.9 %); mp 153-154 °C; MS FAB m/z 325 ($M^+ + H^+$).

Elemental analysis for $C_{16}H_{21}N_2O_3Cl$

Calc'd C, 59.17; H, 6.52; N, 8.62

Found C, 59.08; H, 6.33; N, 8.49

20

(9c) N-[2-(3-Chloro-1H-indol-4-yloxy)-ethyl]-phthalimide as yellowish white crystals: mp 161-163 °C.

Elemental analysis for $C_{18}H_{13}N_2O_3 \cdot 0.33H_2O$

25 Calc'd: C, 62.36; H, 3.97; N, 8.08

Found: C, 62.37; H, 3.68; N, 8.07

30 (9d) N-Benzyl-N-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoro-acetamide as a white solid: (82%); mp 156-158 °C; MS EI m/e 430, 432, 434 (M^+); IR(KBr) 1680 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 3.76 and 3.81 (2t, 2H, $J=1.3$ Hz, $J=1.4$ Hz, rotamers), 4.14 and 4.15 (2t, 2H, $J=1.5$ Hz, $J=1.6$ Hz, rotamers), 4.95 and 4.96 (2s, 2H, rotamers), 6.41 and 6.43 (2d, 1H, $J=8.4$ Hz, $J=8.7$ Hz, rotamers), 7.095 and 7.097 (2d, 1H, $J=8.2$ Hz, $J=8.2$ Hz, rotamers), 7.16 (d, 1H, $J=2.5$ Hz), 7.22-7.41 (m, 5H), 8.27-8.35 (m, 1H).

35

- 16 -

Elemental analysis for $C_{19}H_{15}Cl_2F_3N_2O_2$

Calc'd: C, 52.92; H, 3.51; N, 6.50

Found: C, 52.54; H, 3.26; N, 6.29.

- 5 (9e) N-[2-(3-chloro-1H-indol-4-yloxy)-ethyl]-N-(4-phenyl-butyl)-trifluoroacetamide: (71.4 %), mp 113-114°C; MS *m/e* 438 (M+).

Elemental analysis for $C_{22}H_{22}N_2O_2ClF_3$

Calc'd: C, 60.21; H, 5.05; N, 6.38

- 10 Found: C, 60.51; H, 4.94; N, 6.31.

EXAMPLE 2

12-(3-Chloro-1H-indol-4-yloxy)-ethyl-(4-phenyl-butyl)-amine

15

- A mixture of [2-(3-chloro-1H-indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-trifluoroacetamide (1.15 g, 2.62 mmol) and potassium carbonate (2.53 g, mmol) in a solution of methanol-water (50 mL:3 mL) was heated to reflux for 3 hours. The solvent was removed under vacuum and the crude product was dissolved in methylene chloride (150 mL) and washed with water (100 mL). The aqueous layer was extracted again with methylene chloride (100 mL) and the combined organic layers dried over anhydrous magnesium sulfate, filtered, and the solvent evaporated. The product was purified by flash chromatography (5% methanol-methylene chloride) to afford 847 mg (94.3 %) of a tan oil: MS *m/e* 342 (M+), 344 (M+). The fumarate salt was prepared in isopropanol: mp 195-196 °C.
- 20
- 25

Elemental analysis for $C_{20}H_{23}N_2OCl \cdot 0.5C_4H_4O_4$

Calc'd: C, 65.91; H, 6.29; N, 6.99

Found: C, 66.15; H, 6.38; N, 6.81.

30

This general procedure utilizing N-benzyl-N-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoroacetamide, and N-benzyl-N-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-2,2,2-trifluoroacetamide afforded, respectively:

- 17 -

(2b) Benzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine, 92 %. The fumarate salt was prepared from ethanol as a white powder; mp 201-202 °C; MS EI *m/e* 334, 336, 338 (M⁺).

5 Elemental analysis for C₁₇H₁₆Cl₂N₂O•0.5C₄H₄O₄

Calc'd: C, 58.03; H, 4.61; N, 7.12.

Found: C, 57.88; H, 4.45; N, 6.96.

(2c) 1-[4-(2-Benzylamino-ethoxy)-7-chloro-1H-indol-3-yl]-2,2,2-trifluoro-
10 ethanone: (80 %). The fumarate salt was prepared in ethanol: mp 215 °C (dec); MS
FAB *m/e* 397 (MH⁺).

Elemental analysis for C₁₉H₁₆ClF₃N₂O₂•0.5C₄H₄O₄

Calc'd: C, 55.46; H, 3.99; N, 6.16.

15 Found: C, 55.24; H, 3.80; N, 6.08.

EXAMPLE 3

1-[4-(2-Benzylamino-ethoxy)-1H-indol-3-yl]-2,2,2-trifluoro-ethanone

20

To a solution of N-benzyl-N-[2-(1H-indol-3-(2,2,2-trifluoroethanoyl)-4-yl-oxy)-ethyl]-carbamic acid tert-butyl ester (1.1 g, 2.4 mmol) in methylene chloride (60mL) was added trifluoroacetic acid (TFA) (0.21 g, 1.8 mmol). Thin layer chromatography (TLC) (CH₂Cl₂/CH₃OH, 88/12v/v) showed no change after 1 hour
25 at room temperature. TFA (0.74 g, 6.5 mmol) was added and the mixture stirred 2
hours. Some product was then visible in the TLC. TFA (0.86 g, 7.5 mmol) was added
and the mixture was stirred overnight. TLC showed some starting material. TFA (0.06
g, 0.5 mmol) was added and the mixture was stirred for 1 hour. The reaction mixture
was washed once with saturated NaHCO₃ (30-40 mL). It was dried over MgSO₄.
30 Evaporation of the solvent gave 1.02 g of residue. This was purified by
chromatography on silica gel with a gradient of CH₂Cl₂/CH₃OH (96/4 and 95/5) to
give the product as a light tan oil (0.78 g, 90%).

- 18 -

To a hot solution of fumaric acid (0.2557 g, 2.203 mmol) in EtOH (15 mL) was added a hot solution of the base in EtOH (15 mL). This mixture stood at room temperature for 2 hours. It was filtered to give the title compound as a white powder: 0.5398 g (54%); decomp. >220 °C; MS EI *m/e* 362 (M^+); IR (KBr) 1660 cm^{-1} .

5

Elemental analysis for $\text{C}_{19}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_2 \cdot 0.5\text{C}_4\text{H}_4\text{O}_4$

Calc'd: C, 60.00; H, 4.56; N, 6.66.

Found: C, 60.12; H, 4.40; N, 6.75.

10

This general procedure utilizing benzyl-[2-(1H-indol-4-yloxy)-ethyl]-carbamic acid tert-butyl ester afforded

(3b) N-Benzyl-[2-(1H-indol-4-yloxy)]-ethylamine, (26 %). The fumarate salt was prepared in isopropanol: mp 158-165 °C.

15

Elemental analysis for $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O} \cdot \text{C}_4\text{H}_4\text{O}_4$

Calc'd: C, 65.96; H, 5.80; N, 7.33.

Found: C, 65.94; H, 5.87; N, 7.19.

20

Affinity for the dopamine autoreceptor was established by a modification of the standard experimental test procedure of Seemen and Schaus, European Journal of Pharmacology 203, 105-109 (1991), wherein homogenized rat striatal brain tissue is incubated with ^3H -quinpirole (Quin.) and various concentrations of test compound, filtered and washed and counted in a Betaplate scintillation counter.

25

High affinity for the dopamine D-2 receptor was established by the standard experimental test procedure of Fields, et al., Brain Res., 136, 578 (1977) and Yamamura et al., eds., Neurotransmitter Receptor Binding, Raven Press, N.Y. (1978) wherein homogenized limbic brain tissue is incubated with ^3H -spiroperidol (Spiper.) and various concentrations of test compound, filtered and washed and shaken with Hydrofluor scintillation cocktail (National Diagnostics) and counted in a Packard 460 CD scintillation counter.

30

- 19 -

High affinity for the serotonin 5-HT_{1A} receptor was established by testing the claimed compound's ability to displace [³H] 8-OHDPAT (dipropylaminotetralin) from the 5-HT_{1A} serotonin receptor following the procedure of Hall et al., J. Neurochem. 44, 1685 (1985). This procedure is employed to analogize this property of the claimed compounds with that of buspirone, which is a standard for anxiolytic activity, and, like the compounds of this invention, displays potent affinity for the 5-HT_{1A} serotonin receptor subtype. The anxiolytic activity of buspirone is believed to be, at least partially, due to its 5-HT_{1A} receptor affinity (Vander Maclen et al., Eur. J. Pharmacol. 1986, 129 (1-2) 133-130).

The results of these standard experimental test procedures were as follows:

| Example No. | IC ₅₀ (nM) D ₂ Quin. | IC ₅₀ (nM) D ₂ Spiper | IC ₅₀ (nM) 5-HT _{1A} | Ratio ant/agonist |
|-------------|---|--|---|----------------------|
| (1a) | 10.5 | 184 | 0.71 | 18 |
| (1b) | 14.9 | 183 | 386 | 12 |
| (1c) | 13.6 | 284 | 438 | 21 |
| (1f) | 22.5 | 405 | - | 18 |
| (2a) | 9.55 | 193 | 0.99 | 20 |
| (2b) | 13.6 | 284 | 438 | 21 |
| (2c) | 54.8 | 449 | 102 | 8 |
| (3a) | 5.93 | 286 | 60 | 48 |
| (3b) | 19.4 | 501 | 36 | 26 |

- 20 -

Hence, the compounds of this invention effect the synthesis of the neurotransmitter dopamine and thus are useful in the treatment of dopaminergic disorders such as schizophrenia, Parkinson's disease, Tourette's Syndrome, alcohol
5 addiction, cocaine addiction, and addiction to analagous drugs. These compounds also have affinity for the 5-HT_{1A} receptors and therefore have the ability to modulate serotinerbic activity. As such, they are also useful in the treatment of diseases characterized by disturbances in the serotinerbic systems, such as anxiety, stress, depression, sexual dysfunctions and sleep disturbances.

10

Applicable solid carriers for pharmaceutical compositions containing the compounds of this invention can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintergrating agents or an encapsulating material.
15 In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate,
20 magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient of this invention can be dissolved or
25 suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples
30 of liquid carriers for oral and parenteral administration include water (particularly containing additives as above e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols e.g. glycols) and their derivatives, and oils (e.g. fractionated
35 coconut oil and arachis oil). For parenteral administration the carrier can also be an oily

- 21 -

ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

5 Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Oral administration may be in either liquid or solid composition form.


10 Preferably the pharmaceutical composition is in unit dosage form, e.g. as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged compositions, for example packeted powders, vials, ampoules, prefilled syringes or
15 sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

20 The dosage to be used in the treatment of a specific psychosis must be subjectively determined by the attending physician. The variables involved include the specific psychosis and the size, age and response pattern of the patient.

- 22 -

WHAT IS CLAIMED IS:

(1) A compound of formula I



I

5

in which:

R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cycloalkylalkyl of 6 to 12 carbon atoms, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms, (alkoxyaryl)alkyl of 8 to 12 carbon atoms, thienylmethyl, furanymethyl, pyridinylmethyl, alkylphenyl of 7 to 12 carbon atoms, 4-fluorobutyrophenone or 6-fluoro-1,2-benzisoxazol-yl-propyl;

10

X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

15

Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.

(2) A compound of Claim 1 in which:

20

R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cyclohexylmethyl, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms or (alkoxyaryl)alkyl of 8 to 12 carbon atoms;

X is H, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

25

Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.

- 23 -


- 5 (3) A compound of Claim 1 in which R₁ is alkyl of 1 to 6 carbon atoms, benzyl, halobenzyl, alkoxybenzyl of 8 to 12 carbon atoms or alkylbenzyl of 8 to 12 carbon atoms; X is hydrogen, halogen or trifluoroacetyl and Y is hydrogen or halogen; or a pharmaceutically acceptable salt thereof.
- (4) The compound of Claim 1 which is [2-(1H-indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-amine or a pharmaceutically acceptable salt thereof.
- 10 (5) The compound of Claim 1 which is benzyl-[2-(7-chloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- (6) The compound of Claim 1 which is benzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- 15 (7) The compound of Claim 1 which is 4-fluorobenzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- (8) The compound of Claim 1 which is 4-chlorobenzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- 20 (9) The compound of Claim 1 which is thien-2-ylmethyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- 25 (10) The compound of Claim 1 which is [2-(3-chloro-1H-indol-4-yloxy)-ethyl]-(4-phenyl-butyl)-amine or a pharmaceutically acceptable salt thereof.
- (11) The compound of Claim 1 which is benzyl-[2-(3,7-dichloro-1H-indol-4-yloxy)-ethyl]-amine or a pharmaceutically acceptable salt thereof.
- 30 (12) The compound of Claim 1 which is 1-[4-(2-benzylamino-ethoxy)-7-chloro-1H-indol-3-yl]-2,2,2-trifluoro-ethanone or a pharmaceutically acceptable salt thereof.
- (13) The compound of Claim 1 which is 1-[4-(2-benzylamino-ethoxy)-1H-indol-3-yl]-2,2,2-trifluoro-ethanone or a pharmaceutically acceptable salt thereof.
- 35

- 24 -

(14) The compound of Claim 1 which is (N-benzyl-[2-(1H-indol-4-yloxy)]-ethyl-amine or a pharmaceutically acceptable salt thereof.

(15) A pharmaceutical composition of matter comprising a compound of the formula:

5



I

in which:

10 R_1 is hydrogen, alkyl of 1 to 10 carbon atoms, cycloalkylalkyl of 6 to 12 carbon atoms, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms, (alkoxyaryl)alkyl of 8 to 12 carbon atoms, thienylmethyl, furanylmethyl, pyridinylmethyl, alkylphenyl of 7 to 12 carbon atoms, 4-fluorobutyrophenone or 6-fluoro-1,2-benzisoxazol-yl-propyl;


15

X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

20 Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier therefor.

(16) A method for reducing dopamine synthesis and release in a patient suffering from hyperactivity of the dopaminergic systems, which comprises administering to said patient a compound of the formula:



I

in which:

R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cycloalkylalkyl of 6 to 12 carbon atoms, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms, (alkoxyaryl)alkyl of 8 to 12 carbon atoms, thienylmethyl, furanylmethyl, pyridinylmethyl, alkylphenyl of 7 to 12 carbon atoms, 4-fluorobutyrophenone or 6-fluoro-1,2-benzisoxazol-yl-propyl;

15 X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;


Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

20

or a pharmaceutically acceptable salt thereof, in an amount sufficient to modulate the dopamine systems of the brain.

- 26 -

(17) A method for treating schizophrenia which comprises administering to a patient suffering from schizophrenia, orally or parenterally, a compound of the formula:



I

5

in which:

R₁ is hydrogen, alkyl of 1 to 10 carbon atoms, cycloalkylalkyl of 6 to 12 carbon atoms, arylalkyl of 7 to 12 carbon atoms, (haloaryl)alkyl of 7 to 12 carbon atoms, (alkoxyaryl)alkyl of 8 to 12 carbon atoms, thienylmethyl, furanylmethyl, pyridinylmethyl, alkylphenyl of 7 to 12 carbon atoms, 4-fluorobutyrophenone or 6-fluoro-1,2-benzisoxazol-yl-propyl;

X is hydrogen, halogen, cyano, alkyl of 1 to 6 carbon atoms, acetyl, trifluoroacetyl, trifluoromethyl or formyl;

Y is hydrogen, halogen, alkoxy of 1 to 6 carbon atoms or alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof, in an amount sufficient to alleviate the symptoms of schizophrenia.

INTERNATIONAL SEARCH REPORT

Inte: onal Application No
PCT/US 97/15026

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D209/08 A61K31/40 C07D209/30 C07D409/12 C07D209/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| A | US 5 541 199 A (RICHARD E. MESHAW) 30 July 1996 see example 4 | 1,15 |
| A | --- US 5 013 761 A (EDWARD E. BEEDLE ET AL) 7 May 1991 see column 9 | 1,15 |
| X | --- US 3 906 000 A (JAMES M. MCMANUS) 16 September 1975 cited in the application * column 11, compound on line 44 * | 1 |

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

1 December 1997

Date of mailing of the international search report

09.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 97/15026

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Remark : Although claims 16 and 17 are directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Application No

PCT/US 97/15026

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| US 5541199 A | 30-07-96 | NONE | |
| US 5013761 A | 07-05-91 | US 5420294 A | 30-05-95 |
| | | US 5554640 A | 10-09-96 |
| | | AT 117980 T | 15-02-95 |
| | | AU 609997 B | 09-05-91 |
| | | AU 3601989 A | 07-12-89 |
| | | CA 1336911 A | 05-09-95 |
| | | CN 1045774 A,B | 03-10-90 |
| | | DE 68920913 D | 16-03-95 |
| | | DE 68920913 T | 22-06-95 |
| | | DK 271889 A | 05-12-89 |
| | | EP 0345056 A | 06-12-89 |
| | | ES 2067541 T | 01-04-95 |
| | | IE 66225 B | 13-12-95 |
| | | IL 90498 A | 08-12-95 |
| | | JP 2025416 A | 26-01-90 |
| | | PT 90716 B | 01-03-95 |
| | | SU 1795965 A | 15-02-93 |
| US 3906000 A | 16-09-75 | US 3833591 A | 03-09-74 |
| | | BE 795451 A | 16-08-73 |
| | | DE 2306605 A | 06-09-73 |
| | | FR 2181738 A | 07-12-73 |
| | | GB 1418354 A | 17-12-75 |
| | | JP 1027972 C | 25-12-80 |
| | | JP 48092400 A | 30-11-73 |
| | | JP 55016434 B | 01-05-80 |
| | | US 3904645 A | 09-09-75 |
| | | US 3898245 A | 05-08-75 |